



Addition to carbon-hetero-multiple bonds. Ist Lect.

(P/KS)

These reactions involve addition to C=O, C=N and C=S double bonds and C≡N triple bond. Since C=O, C=N and C≡N bonds are strongly polar with the carbon always the positive end, there is never any doubt about orientation of unsymmetrical addition to these bonds. Nucleophiles always go to the carbon while electrophilic attacking species go to oxygen or nitrogen. Free radical addition to carbon-hetero bonds are rare. In most of the cases, nucleophile attacks first and these reactions are regarded as nucleophilic addition.




When A and B are H, R or Ar the substrate is an aldehyde or ketone and they almost never ^{under}go substitution, due to the extremely poor nature of H, R and Ar as leaving group. For acids and their derivatives (B = OH, OR, NH₂, etc.) addition is seldom found, because these are much better leaving groups.

Reaction with hydride donors


Many reactions of aldehydes and ketones involve transfer of hydride ion (H^{\ominus}) from donor molecule to carbonyl carbon. The common hydride donors are - lithium aluminium hydride, sodium borohydride and aluminium alkoxides, while in Cannizzaro's reaction hydride ion transfer occurs from an already attacked aldehyde molecule.

(1) lithium Aluminium Hydride LiAlH₄

It is a reagent widely used for reduction of carbonyl compound to alcohols. All the four hydrogen atoms are available to the carbonyl compound as hydride ions, thus converting it into a metal complex, which then gets decomposed by acetone.






The reaction appears to be a concerted process in which the hydride ion attacks carbonyl carbon and simultaneously the carbonyl oxygen starts complexing with aluminium ion.




Reduction of Carboxylic acids and its derivative.




Reduction of carboxylic acids and its derivatives, such as esters, acid anhydrides and acier chlorides result in the formation of corresponding alcohols. The order of reactivity of carboxylic acid and its derivatives toward LiAlH_4



Carboxylic acid reacts with LiAlH_4 to form carboxylate anion, and this is followed by the addition of hydride ion to the carbonyl group to give tetrahedral intermediate, which is further reduced to primary alcohol.




Similarly esters can be reduced.




(3)

benzyl substituted amides give primary amines, whereas, N-sub amide gives secondary amines and N,N-disubstituted amide give tertiary amines. Reduction of amide with LiAlH₄ gives iminium ion, which is more electrophilic than starting amide so it gets reduced to amino.




Reduction of nitriles to amines



NaBH_4 does not generally reduce nitriles, but does so in acidic solvents when a CoCl_2 catalyst is added or in presence of Raney Ni. The reaction is of third scope. It is not possible to stop with the addition of only 1 mole of hydrogen.


N-alkyl nitinium ions are reduced to secondary amines by NaBH_4 .




④ NaBH_4

NaBH_4 is a mild reducing agent, which reacts rapidly with aldehydes and ketones but only slowly with esters. Neither NaBH_4 nor LiAlH_4 react with isolated C=C double bond. The alkali metal hydrides such as NaH are unsuitable reducing agents because of their insolubility in organic solvents and their powerful effect as catalyst for base catalysed condensation.


The most commonly used reagent for reduction of aldehydes and ketones is Sodium borohydride NaBH_4 . Since BH_4^- contains four hydride ions it is capable of reducing four molecules of aldehydes or ketones. Reaction proceeds by successive transfer of hydride ions from the boron to four different carbonyl carbons. As all the four hydrides are transferred there are actually several reducing species which function during course of reduction.



Reduction is carried out either in alcohol or in water. While reductions with LiAlH₄ are done in a protic solvent e.g. anhyd. ether.



(3) Aluminium alkoxides: This reaction involves the transfer of hydride ions from the metal salts of primary and secondary alcohols to the carbonyl carbon. The reduction of carbonyl compounds to alcohols with Aluminium isopropoxide is very important reaction known as Meerwein-Ponndorf-Verley Reduction. Isopropanol is used as the solvent. Acetone is distilled off as soon as it is formed from the reaction mixture.



(4) Cannizzaro's Reaction

Aldehydes which do not have α -hydrogen atoms cannot undergo base-catalysed condensation, they react with bases by disproportionation involving the transfer of hydride ions. It is self oxidation-reduction reaction. Crossed Cannizzaro's reaction between one molecule of such aldehyde and another molecule of formaldehyde leads to reduction of the former and the oxidation of the latter.

